

Official Translator - Interpreter Accredited by the
Colombian Ministry of Justice with Resolution No. 16-37.

Traductora - Intérprete Oficial Acreditada con Resolución
No. 16-37 del Ministerio de Justicia de Colombia.

Rene Camacho Cordoba, MD
Medical Register No. 13-844/95
Colombian Health Ministry

Cartagena de Indias, Colombia
August 16, 1999

Patient: Joseph Hertrich
Nationality: USA
Passport No. 87728840
Sex: Masculine
Age: 43 years old
Laterality: Right-handed
Personal antecedent: Gall bladder surgery two years ago.

Should be "bladder surgery"

Reason for consultation:

I was asked to visit this patient for a clinical evaluation due to a probable infection of undetermined cause and origin. The Clinical History (CH) of this patient is extensive and complex. The patient presents numerous medical records and blood tests since the year 92 up to the present. This require a detailed study and later consultation with a specialist in Hematology, a Neurosurgeon in the City of Cartagena de Indias, Colombia.

Physical observation of the Patient:

Forty-three-year-old male adult, oriented in respect to time and location. Moderate to severe stress. No speech alterations. Frequent involuntary spasm in the lower extremes. He states he feels pains in the lower and upper extremes. Thin. He needs an orthopedic apparatus (walker) to be able to move around.

Clinical evaluation:

Once the Clinical History and past blood tests of the patient were revised, with the cooperation efforts of the specialist in Hematology, a tentative conclusion was reached. The blood tests clearly show Leukocytes and Neutrophilia. This indicates that the patient, Joseph Hertrich, presents an infectious process which was left with an efficient treatment for a period of seven years.

Due to the great deterioration of the patient's health, his slight improvement in the past with the use of oral antibiotics and the indications in the blood tests or complete blood count (CBC), it was determined that the best and prudent course of action would be to start of treatment with antibiotics intravenously immediately. The initial medicine chosen was Rocefin 1 gram twice a day.